

# HEALTH CARE POLICY

---

|                        |                |
|------------------------|----------------|
| Date of policy review: | September 2026 |
| Date for review:       | September 2027 |
| Lead for review:       | School Nurse   |

## Contents

|                                                                      |    |
|----------------------------------------------------------------------|----|
| Contents.....                                                        | 2  |
| Preface .....                                                        | 3  |
| Aim .....                                                            | 3  |
| Confidentiality and Information Sharing .....                        | 3  |
| Documentation .....                                                  | 4  |
| Working with Parents/Guardians .....                                 | 4  |
| Medical Services and Support.....                                    | 4  |
| Reporting Sickness .....                                             | 5  |
| Medication Management .....                                          | 5  |
| Prescribed Medicines.....                                            | 6  |
| Emergency Medical Procedures .....                                   | 6  |
| Health Care Plans .....                                              | 7  |
| Related Policies .....                                               | 8  |
| Appendices.....                                                      | 8  |
| Appendix 1: Health Plan Template .....                               | 8  |
| Appendix 2: Protocol for Homely remedies for Guildhouse.....         | 10 |
| Appendix 3: FLOWCHART OF KEY QUESTIONS FOR INFORMATION SHARING ..... | 17 |
| Appendix 4: GOOD HYGIENE PRACTICE .....                              | 18 |
| Appendix 5: Helpful organisation contacts.....                       | 19 |
| Appendix 6: PREMIER HOUSE MEDICAL EMERGENCY FLOW CHART .....         | 21 |

## Preface

Guildhouse School is committed to ensuring that all students receive appropriate medical care during their time with us. This policy provides clear guidelines in the following areas:

- Medical care offered by the school
- Confidentiality and information sharing
- Documentation
- Collaboration with parents/guardians
- Procedures for reporting sickness
- Administration, record-keeping, and disposal of medicines
- Health care plans

This policy applies to Guildhouse School in all contexts, including its boarding provision. It is informed by the following statutory and best-practice guidance:

National Minimum Standards for Boarding Schools (NMS 2022) – particularly Standard 7: Boarders’ health and wellbeing, which outlines expectations for medical care, access to healthcare professionals, and medication management in residential settings.

Equality Act 2010 – requiring schools to make reasonable adjustments to ensure pupils and staff with disabilities are not disadvantaged in accessing healthcare.

Guildhouse School is committed to ensuring that healthcare provision is consistent, inclusive, and responsive to the needs of all pupils, including those in boarding, in line with safeguarding and health standards.

## Aim

To collaborate with parents, guardians, students, and healthcare professionals to ensure students requiring medical treatment can do so in a safe and supportive environment, enabling them to continue their education effectively.

## Confidentiality and Information Sharing

Guildhouse School is committed to handling all personal and medical information in accordance with the UK General Data Protection Regulation (UK GDPR) and the Data Protection Act 2018 (DPA 2018). In line with data protection regulations, students under 18 require parental or authorised representative consent for the school to process their health information. The school nurse is bound by professional confidentiality but may share medical information when:

- It is in the best interest of the student or the wider school community, even if consent cannot be obtained.
- An emergency arises where the student is unable to give consent.

Information shared will follow the guidelines in Appendix 3 (Information Sharing Flowchart). The reasons and details of shared information must be documented as per school procedures.

[https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment\\_data/file/721581/Information\\_sharing\\_advice\\_practitioners\\_safeguarding\\_services.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/721581/Information_sharing_advice_practitioners_safeguarding_services.pdf)

In safeguarding contexts, the school follows statutory guidance from:

Working Together to Safeguard Children (2026)

Keeping Children Safe in Education (KCSIE 2026)

## Documentation

Medical interactions and communications are recorded securely in the school's designated systems (e.g., iSAMs) and stored in secure personal files on OneDrive/SharePoint. Hard copies are returned to the student or disposed of as confidential waste. Access is limited to authorised staff only.

New students go through a medical-health arrival questionnaire upon joining with the Nurse, detailing their medical history, current treatments, allergies, immunisations, mental health, and eating habits.

## Working with Parents/Guardians

Guildhouse School values collaboration with parents, guardians, or agents to address student health needs effectively. Parents are expected to provide accurate and up-to-date information about their child's medical conditions, medications, and any special requirements. This includes:

- Medicine name, dosage, and administration method
- Frequency and timing
- Side effects impacting behaviour or performance
- Changes in medication

The school will respect cultural and religious considerations while ensuring health and safety.

## Medical Services and Support

Students are directed to register with a local GP upon arrival. The school provides:

- Ongoing support from the School Nurse. There is a clear procedure for seeing the Nurse, usually by contacting them on Microsoft Teams. If a situation occurs during a lesson, the teacher contacts the Nurse to see if they can attend to the student.
- In the case of the Nurse being off-site, students are directed to Reception who will then liaise in the first instance and request for First Aid trained Student Services/Administration staff to see the student.
- If they are not available, reception will liaise with the Academic and Pastoral Hub (Middle/Senior Leaders, who are First Aid Trained, for someone to see the student.
- Support for appointments with GPs, dentists, and optometrists
- Staff accompaniment for students under 18 during medical appointments (unless parental consent is given for unaccompanied visits)
- Students over 18 can attend medical appointments unaccompanied but it must be confirmed they wish to do this; some may request for a member of staff to attend with them

- Support for students with chronic conditions such as asthma, diabetes, epilepsy, and allergies, including relevant staff training and emergency action plans
- Mental health counselling services arranged through the School Nurse with the School Counsellor – these may sometimes be advised
- School Staff/Boarding Staff should consult with the National Health Service (NHS) helpline 111 for non-emergency medical advice and assistance.

Individual cases are assessed to determine the level of support required. Where resources are insufficient to meet a student's needs, alternative arrangements will be discussed with parents or guardians.

## Reporting Sickness

Students feeling unwell must follow the attendance procedures outlined in the Student and Boarding Handbook. After assessment, the following actions may be taken:

1. Return to class if deemed fit.
2. Rest in their residence or the boarding medical bay, with staff informed.
3. Arrange a GP or private doctor appointment, with parental consent required for private consultations.
4. Direct the student to trained first aid staff if the School Nurse is unavailable by directing them to reception.

For illnesses lasting more than three days, proof of sickness via an NHS doctor's certificate may be required and the school reserves the right to insist that a student attends an NHS Doctor to enable proof to be received.

Sickness absences need to be reported to the school nurse (working with SLT) in line with attendance policy in order for the absence to be authorised.

## Medication Management

It is important that students who need to take medication at school are involved as closely as possible in the arrangements made for them. When planning for medical care at school the following is taken into consideration:

- Independent management of needs
- Supervised administration of medication
- Staff administration of medication

### Storage

- Medicines are kept in locked cupboards or refrigerators in medical rooms. Members of staff who use the refrigerator must be made aware of the importance of keeping medicine safe and secure.
- Students competent to self-medicate (Gillick Competent) may store both over the counter medication and prescribed medication in personal safes.
- Students will carry their own inhalers/EpiPens with them.
- Spare emergency inhalers/EpiPens are kept in medical rooms and accommodation, clearly labelled.
- Advice on the storage of medicines will be sought from a qualified pharmacist when required.

- Medicines may be potentially harmful to anyone for whom they are not prescribed. The school acknowledges that it has a duty to ensure that risks to the health of others are properly controlled.
- All staff should be familiar with the normal procedures for avoiding infection and will follow the basic hygiene procedures detailed in the Infection Control Guidelines issued in the UKHSA 2022 (Health protection in children and young people settings). (See Appendix 4.)
- Medication should be taken to school only when it is needed. Often medication can be prescribed in dose frequencies, which enable it to be taken outside school hours.

## **Administration**

- Medicine administration is supervised by the Nurse or authorised staff.
- Details are recorded on iSAMS, including dosage & time.
- Risk assessments and health plans are implemented for controlled medications.
- Details will be completed on iSAMS for each student receiving medication.
- The health plan of medication administration will be reviewed with the student and nurse, at set intervals, to ensure the student's medical needs are being met.
- Any changes to the plan will be updated on iSAMS and the appropriate staff informed by the School Nurse.

## **Non-Prescribed Medicines**

- Staff must check for previous doses, allergies, or contraindications before administering homely remedies.
- Staff must not compel a student to take medication.
- Parental consent for under-16s is required, unless the student is assessed as Gillick Competent.

## **Disposal**

- Medicines are disposed of via local pharmacies and recorded in a disposal log. Sharps are placed in designated boxes.

## **Prescribed Medicines**

Any member of staff giving medicines to a student should observe the following procedure in cooperation with a colleague

- confirm the student's name agrees with that on the medication
- check the written instructions provided by the parents or doctor
- confirm the prescribed dose
- check the expiry date
- check notes on iSAMS

Staff record on iSAMS each time they give medication to a student and alert the note to the school Nurse.

## **Emergency Medical Procedures**

The Nurse and Deputy Head Teacher Pastoral will ensure that all staff know how to call the Emergency Services. Names of staff qualified to administer First Aid will be posted in rooms around the school and in the Boarding residence.

In medical emergencies:

- Staff must call 999/112 and provide detailed information.
- A member of staff will accompany the student to the hospital and act in loco parentis.
- In the case of a student self-referring, a member of staff would be assigned to accompany and stay until it's decided by SLT that the student can remain on their own.
- Parents or guardians will be informed promptly.
- If a student is admitted, the school will make arrangements for an overnight bag to be brought to the hospital and find out if it is appropriate for friends to visit.
- During a hospital stay, the school will liaise regularly with medical staff and ensure appropriate accompaniment and visits are organised.
- Medication currently being taken by the student should be brought to the hospital.
- Refusal of medication is documented, and emergency services are contacted if necessary.
- If a student is taken to hospital during school hours, the Head Teacher/SLT and School Nurse must be informed.
- The school will then undertake to inform the parents/guardian/agent and keep the parents/guardian/agent updated.
- A member of staff will be with the student to facilitate discharge from the hospital.
- If a student is taken to hospital during boarding hours (Monday – Friday 1730-0830 / all day Saturday and Sunday) then call the emergency telephone number. The on-call senior member of staff will inform the parents/guardian/agent as soon as is practical.
- When a student is taken to hospital by a member of staff, they should also take with them all medication the student is currently taking. If a student refuses to take medication, the school will record this on Shackleton and inform the student's parents.

## Sexual Health

All students at Guildhouse school will be taught about sexual health as part of the PSHE/RSE programme. Students can liaise with the School Nurse confidentially if they have any concerns or questions regarding sexual health. Students will be directed onto local services where appropriate. Information of local services that provide free testing and protection will be on notice boards around the school. There are multiple clinics within walking distance including The Brook, CNWL & 56 Dean Street (LGBTQ+).

## Health Care Plans

Individual health care plans outline the required level of support for students with medical needs. Plans are developed with input from:

- The student (if appropriate)
- Parents/guardians/agents
- The school nurse and relevant staff
- External healthcare professionals

Plans include medication details, emergency procedures, and review intervals. They ensure tailored support and effective communication among all stakeholders.

Health Care Plans will be reviewed annually, or sooner if there is a change in the student's medical condition or care needs. Guildhouse School is committed to ensuring that all healthcare arrangements are inclusive and accessible. In line with the Equality Act 2010, reasonable adjustments will be made to support students with disabilities, ensuring equitable access to medical care and support.

## Monitoring and Review

This policy is monitored and reviewed annually or sooner if triggered by changes in legislation, statutory guidance or significant incidents by the Deputy Head Pastoral and School Nurse. Any updates are communicated to relevant staff as part of training, and stakeholders

## Related Policies

Health & Safety Policy

First Aid Policy

Attendance Policy

## Appendices

### Appendix 1: Health Plan Template

|                                                               |  |
|---------------------------------------------------------------|--|
| Student Name:                                                 |  |
| D.O.B.:                                                       |  |
| Gender:                                                       |  |
| Residence:                                                    |  |
| Names Health Coordinator:                                     |  |
| Family Contact on Shackleton:                                 |  |
| Medical Diagnosis/Condition<br>(include medicine prescribed): |  |

|                               |  |
|-------------------------------|--|
| Medical Professional Contact: |  |
|-------------------------------|--|

|                   |  |
|-------------------|--|
| Name:             |  |
| Telephone Number: |  |
| Address:          |  |

| Assessment    |          |
|---------------|----------|
| Medical Needs | Symptoms |
|               |          |
|               |          |

|                                      |  |
|--------------------------------------|--|
| Daily Health Plan to be implemented: |  |
| Health Plan Implemented by whom:     |  |

|                                                  |  |
|--------------------------------------------------|--|
| Potential Complications and Emergency Situation: |  |
| Agreed Actions to take:                          |  |
| Actions Implemented by whom:                     |  |

|                    |  |
|--------------------|--|
| Review Due:        |  |
| Review By:         |  |
| Date Reviewed:     |  |
| Outcome of Review: |  |

|                |  |
|----------------|--|
|                |  |
| Date of Plan:  |  |
| Review Date:   |  |
| Date Reviewed: |  |
| Reviewed by:   |  |

## Appendix 2: Protocol for Homely remedies for Guildhouse

### Guidance

This is a list of medication that can be administered to students that have not been prescribed by a doctor. The aim of these guidelines is to provide safe treatments for commonly presented conditions. This policy does not supersede the need to contact a doctor, by any member of staff, if they are unsure or there is any doubt about the condition being treated.

Administration of these remedies should be given in accordance with the patient information leaflets provided in each pack, taking into account that there are no contraindications or previous allergies to the medication.

Any medication administered must be clearly recorded on the medical section in Shackleton and the school Nurse informed of the time and dose that the medication was given.

### Homely Remedies

| CONDITION                               | TREATMENTS                   |
|-----------------------------------------|------------------------------|
| Cold and 'flu like symptoms'            | Paracetamol or decongestant  |
| Muscular Pain Relief, anti-inflammatory | Paracetamol, Ibuprofen       |
| Allergy relief, hay fever               | Allergy and Hay fever relief |
| Diarrhoea                               | Loperamide hydrochloride     |
| Indigestion relief                      | Gaviscon or Rennie           |
| Rehydration                             | Rehydration treatment        |

|                       |          |
|-----------------------|----------|
| Minor cuts and grazes | Plasters |
|-----------------------|----------|

## Paracetamol

### Indications

|                     |                                                            |
|---------------------|------------------------------------------------------------|
| When it can be used | Pain relief for mild to moderate pain, pyrexia (fever)     |
| Do Not Give         | In conjunction with other medicines containing paracetamol |

### Treatment to be given

|                           |                                                                     |
|---------------------------|---------------------------------------------------------------------|
| Name of Medicine          | Paracetamol 500mg                                                   |
| Dose                      | 1 to 2 tablets                                                      |
| Route                     | Oral                                                                |
| Frequency                 | Four to Six hours between doses                                     |
| Max dose in 24 hrs        | 4g (8 tablets)                                                      |
| Follow up                 | Inform GP if symptoms persist                                       |
| Warning/Adverse reactions | Side effects rare – rash, blood disorders, liver damage in overdose |

## Ibuprofen

### Indications

|                     |                                                                                                                                                                                                                                                                                                                                                                                                         |
|---------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| When it can be used | Pain relief for mild to moderate pain, migraine, musculoskeletal pain.                                                                                                                                                                                                                                                                                                                                  |
| Do not give         | Asthma, pregnancy, known hypersensitivity to aspirin, ibuprofen or other NSAID. Current or previous history of dyspepsia or peptic ulceration, patients taking oral anticoagulants, warfarin, heparin, aspirin or other NSAIDs, patients taking lithium, methotrexate, tacrolimus, ciclosporin, and patients with known severe cardiac disease, heart failure, oedema, hypertension or renal impairment |

### Treatment to be given

|                            |                                                                                                                 |
|----------------------------|-----------------------------------------------------------------------------------------------------------------|
| Name of Medicine           | Ibuprofen 200 mg tablets                                                                                        |
| Dose                       | 200mg – 400mg                                                                                                   |
| Route                      | Oral                                                                                                            |
| Frequency                  | Every 8 hours                                                                                                   |
| Max Dose in 24 hrs         | 6 x 200mg tablets                                                                                               |
| Follow up                  | If condition worsens or symptoms persist then<br>Seek further medical advice                                    |
| Warnings/Adverse Reactions | Discontinue if indigestion or other gastro – intestinal symptoms develop e.g. haematemesis (blood in vomit)     |
| Advice to student          | Take Medicine with or after food or milk. Ibuprofen may be taken with paracetamol if necessary. Advise Students |

|  |                                                                                                      |
|--|------------------------------------------------------------------------------------------------------|
|  | not to take another non-steroidal anti – inflammatory (NSAIDS) containing products at the same time. |
|--|------------------------------------------------------------------------------------------------------|

## Decongestant

### Indications

|                     |                                                                                                                                                                                                                                                                                                                              |
|---------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| When it can be used | For relief of colds and flu symptoms                                                                                                                                                                                                                                                                                         |
| Do not give         | If allergic to paracetamol, ascorbic acid, or any of the ingredients contained within this product.<br>Have kidney or liver problems, overactive thyroid, diabetes, high bp or heart disease Taking drugs for heart problems or tricyclic antidepressants or have been prescribed drugs for depression in the last two weeks |

### Treatment to be given

|                            |                                                 |
|----------------------------|-------------------------------------------------|
| Name of medicine           | Decongestant                                    |
| Dose                       | 1 sachet every 4-6 hrs                          |
| Route                      | Oral                                            |
| Frequency                  | One sachet every 4-6 hrs                        |
| Max Dose in 24hrs          | No more than 6 sachets                          |
| Follow up                  | See GP if condition worsens or does not improve |
| Warnings/Adverse Reactions | Rare – allergic reactions, such as skin rash    |

## Hayfever and allergy relief

### Indications

|                     |                                                           |
|---------------------|-----------------------------------------------------------|
| When it can be used | Symptomatic relief of allergy such as hayfever, urticaria |
| Do Not Give         | Kidney problems, pregnant or breastfeeding                |

### Treatment to be given

|                            |                                                                                     |
|----------------------------|-------------------------------------------------------------------------------------|
| Name of Medicine           | Hay fever and allergy relief tablets                                                |
| Dose                       | One tablet daily                                                                    |
| Route                      | Oral                                                                                |
| Frequency                  | Once daily                                                                          |
| Max dose in 24hrs          | 1                                                                                   |
| Follow up                  | If no relief, refer to GP                                                           |
| Warnings/Adverse Reactions | Rare – headache, dizziness, dry mouth, drowsiness, stomach or intestinal discomfort |

## Cough Linctus

### Indications

|                     |                                                 |
|---------------------|-------------------------------------------------|
| When it can be used | For relief of chesty coughs                     |
| Do not give         | Fructose intolerance or allergic to ingredients |

### Treatment to be given

|                            |                                                    |
|----------------------------|----------------------------------------------------|
| Name of medicine           | Cough Linctus                                      |
| Dose                       | 10ml (2x5ml spoonful)                              |
| Frequency                  | 2-3 hourly                                         |
| Max dose in 24hrs          |                                                    |
| Follow up                  | Refer to GP if symptoms persist with a temperature |
| Warnings/Adverse Reactions | Rare – stomach upset                               |

## Loperamide hydrochloride

### Indications

|                     |                                                                                                                                                                                                                                                                                                        |
|---------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| When it can be used | To relieve symptoms of diarrhoea                                                                                                                                                                                                                                                                       |
| Do not use          | Cases of severe diarrhoea after taking antibiotics, inflammatory bowel conditions such as ulcerative colitis, constipation, acute dysentery.<br>Advice needed if patient has AIDS and stomach becomes swollen.<br>If diarrhoea lasts more than two weeks (or is related to IBS) if diarrhoea is severe |

### Treatment to be given

|                            |                                                                                                                                                                                                                                                  |
|----------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Name of Medicine           | Loperamide Hydrochloride                                                                                                                                                                                                                         |
| Dose                       | Two tablets initially, followed by 1 capsule after each loose bowel movement                                                                                                                                                                     |
| Route                      | Oral                                                                                                                                                                                                                                             |
| Frequency                  | Take after each loose bowel movement                                                                                                                                                                                                             |
| Max dose in 24hrs          | 16mg daily                                                                                                                                                                                                                                       |
| Follow up                  | If symptoms persist after 3 days, seek medical help                                                                                                                                                                                              |
| Warnings/Adverse Reactions | Abdominal cramps, nausea, vomiting, tiredness, drowsiness, dizziness, dry mouth and skin reactions                                                                                                                                               |
| Advice to student          | Drink plenty of fluids to maintain hydration. To prevent spread of infection wash hands after going to the toilet, before you touch food. Rehydration therapy may be needed in cases of severe diarrhoea where large amounts of fluids are lost. |

## Toothache tincture

### Indications

|                      |                                                                                                                                                                                                                                                                                                                                                   |
|----------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| When it can be given | To relieve sore gums                                                                                                                                                                                                                                                                                                                              |
| DO NOT GIVE          | <p>If suffering from stomach ulcer. Allergic to any ingredients contained within the product.</p> <p>DO NOT GIVE to children or adolescents under the age of 16. There is a junior Bonjela available for this age group or boots own sore gum</p> <p>Seek advice if pregnant or breastfeeding. Linked to Reyes syndrome if given to children.</p> |

### Treatment to be given

|                            |                                                                |
|----------------------------|----------------------------------------------------------------|
| Name of Medicine           | Tooth Tincture                                                 |
| Dose                       | One centimetre applied to sore area                            |
| Route                      | Oral                                                           |
| Frequency                  | Every 3hrs                                                     |
| Max Dose in 24 hrs         |                                                                |
| Warnings/Adverse reactions | Possible allergic reaction, symptoms may like those of asthma. |
| Advice                     | If symptoms persist after 7 days, seek advice                  |

## Indigestion Remedies

### Indications

|                     |                                                                                                                                                                                   |
|---------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| When it can be used | To relieve Indigestion                                                                                                                                                            |
| DO NOT GIVE         | <p>Within two hours of taking other medicine by mouth as it may be less effective. Seek advice if suffering from phenylketonuria as this product is sweetened with aspartine.</p> |

### Treatment to be given

|                            |                                                                                                                                   |
|----------------------------|-----------------------------------------------------------------------------------------------------------------------------------|
| Name of Medicine           | Indigestion Remedy                                                                                                                |
| Dose                       | 2-4 tablets after meals and at bedtime                                                                                            |
| Route                      | Oral                                                                                                                              |
| Frequency                  | After meals                                                                                                                       |
| Max dose in 24 hrs         |                                                                                                                                   |
| Warnings/Adverse reactions | Too many tablets can cause bloating. Rare- allergic reaction                                                                      |
| Advice to students         | Look at lifestyle of student if appropriate. Do they smoke, drink too much coffee, alcohol or eat too many fatty food, chocolate. |

## Rehydration

### Indications

|                     |                                                                                                                                                                       |
|---------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| When it can be used | To replace salts after diarrhea or excessive vomiting                                                                                                                 |
| DO NOT USE          | Ask Doctor or pharmacist if suffering from intestinal obstruction, inflammatory bowel disease, diabetes, kidney or liver failure or on a low potassium or sodium diet |

### Treatment to be given

|                            |                                                                                     |
|----------------------------|-------------------------------------------------------------------------------------|
| Name of medicine           | Rehydration treatment                                                               |
| Dose                       | 1 – 2 sachets with 200ml of water after each loose motion                           |
| Route                      | Oral                                                                                |
| Max Dose in 24 hrs         | 6                                                                                   |
| Warnings/Adverse reactions | Rare- Allergic reaction                                                             |
| Follow Up                  | If symptoms persist, medical advice must be sought                                  |
| Advice to Student          | If vomiting is present the solution should be given in small frequent doses in sips |

## Calamine Lotion


### Indications

|                     |                                          |
|---------------------|------------------------------------------|
| When it can be used | To relieve sun, burn and skin irritation |
| DO NOT USE          | If allergic to lotion                    |

### Treatment to be given

|                  |                                      |
|------------------|--------------------------------------|
| Name of Medicine | Calamine Lotion                      |
| Dose             | Apply as necessary to affected area. |
| Frequency        | As required                          |
| Follow up        | See GP if symptoms persist           |

## Appendix 3: FLOWCHART OF KEY QUESTIONS FOR INFORMATION SHARING



## Appendix 4: Good Hygiene Practice

For more advice contact your local Health Protection Unit or school health service.

### Handwashing

It is one of the most important ways of controlling the spread of infections, especially those that cause diarrhoea and vomiting and respiratory disease. The recommended method is the use of liquid soap, water and paper towels. Always wash hands after using the toilet, before eating or handling food, and after handling animals. Cover all cuts and abrasions with waterproof dressings.

### Coughing and sneezing easily spread infections

Children and adults should be encouraged to cover their mouth and nose with a tissue. Wash your hands after using or disposing of tissues. Spitting should be discouraged.

### Cleaning of the environment

Including toys and equipment should be frequent, thorough, and follow national guidance e.g. use colour coded equipment, COSHH, correct decontamination of cleaning equipment. Monitor cleaning contracts and ensure cleaners are appropriately trained with access to Personal Protective Equipment PPE (see below).

### Cleaning of blood and body fluid spillages

All spillages of blood, faeces, saliva, vomit, nasal, and eye discharges should be cleaned up immediately (always wear PPE). When spillages occur, clean using a product which combines both a detergent and a disinfectant. Use as per manufacturer's instructions and ensure it is effective against bacteria and viruses, and suitable for use on the affected surface. NEVER USE mops for cleaning up blood and body fluid spillages use disposable paper towels and discard clinical waste as described below. A spillage kit should be available for blood spills.

### Personal Protective Clothing (PPE)

Disposable non powdered vinyl or latex free CE marked gloves and disposable plastic aprons, must be worn where there is a risk of splashing or contamination with blood/ body fluids. Goggles should also be available for use if there is a risk of splashing to the face. Correct PPE should be used when handling cleaning chemicals.

### Laundry

It should be dealt with in a separate dedicated facility. Soiled linen should be washed separately at the hottest wash fabric will tolerate. Wear PPE when handling soiled linen. Soiled children's clothing should be bagged to go home, never rinse by hand.

### Clinical waste

Always segregate domestic and clinical waste in accordance with local policy. Used sanitary products, gloves, aprons and soiled dressings should be stored in correct clinical waste bags in foot operated bins. All clinical waste must be removed by a registered waste contractor. All clinical waste bags should be less than 2/3rds full and stored in a dedicated, secure area whilst awaiting collection.

### Sharps injuries and bites

If skin is broken make wound bleed/wash thoroughly using soap and water. Contact GP or go to Accident and Emergency immediately.

## **Appendix 5: Helpful organisation contacts**

### **Allergy UK**

Allergy Help Line: 01322 619898

Website: [Allergy UK | National Charity](#)

### **The Anaphylaxis Campaign**

Helpline: (01252) 542029

Website: [www.anaphylaxis.org.uk](http://www.anaphylaxis.org.uk)

### **Asthma UK (formerly the National Asthma Campaign)**

Adviceline 0300 222 5800 (Mon-Fri 9am to 5pm)

Website: [www.asthma.org.uk](http://www.asthma.org.uk)

### **Council for Disabled Children**

Website: [Council for Disabled Children](#)

### **Contact a Family**

Helpline: 0808 808 3555

Website: [Contact: the charity for families with disabled children](#)

### **Diabetes UK**

Careline: 0345 123 2399 (Weekdays 9am to 5pm)

Website: [Diabetes UK - Know diabetes. Fight diabetes. | Diabetes UK](#)

### **Department for Education**

Website: [Department for Education - GOV.UK](#)

### **Department of Health & Social Care**

Tel: (020) 7210 4850

Website: [Department of Health and Social Care - GOV.UK](#)

### **Equality and Human Rights Commission | Creating a fairer Britain**

[Homepage](#) | [EHRC](#)

### **Equality Advisory and Support Service (EASS)**

Telephone: 0808 800 0082

Website: [Equality Advisory and Support Service](#)

### **Epilepsy Action**

Freephone Helpline: 0808 800 5050 (Monday – Thursday 9am to 4.30pm, Friday 9am to 4pm)

Website: [Home - Epilepsy Action](#)

### **Health and Safety Executive (HSE)**

HSE Infoline: 0300 790 6787 (Mon-Fri 8am-6pm)

Website: [HSE: Information about health and safety at work](#)

### **MENCAP**

Telephone: (020) 7454 0454

Website: [Learning Disability - Down syndrome - Williams syndrome | Mencap](#)

### **National Eczema Society**

Helpline: 020 7281 3553


Website: [Home - National Eczema Society](#)

### **Psoriasis Association**

Tel: 01604 251 620

Website: [The Psoriasis Association](#)

## Appendix 6: PREMIER HOUSE MEDICAL EMERGENCY FLOW CHART





---

GUILDHOUSE SCHOOL  
LONDON

---



Creating tomorrow's change makers.